


# Inflammatory Bowel Disease

## Normal Colon

Crypts should be oriented parallel to one another, perpendicular to the surface (like test tubes in a rack), resting on the muscularis mucosae.



### Regional Variation

Right Colon	Left Colon
More lymphocytes	Less lymphocytes
Paneth cells normal	Paneth Cells abnormal
Fewer goblet cells	More goblet cells

Some architectural distortion and muciphages in the rectum is considered normal.  
 Intraepithelial lymphocytes (and even rare neutrophils) over lymphoid follicles is also normal.

## Patterns of Damage in IBD

The inflammation in IBD is characterized by the presence/absence of “Activity,” defined as neutrophilic inflammation of the epithelium and lamina propria, and “Chronicity,” including architectural distortion, a basal lymphoplasmacytosis, and Paneth cell metaplasia.

These words are combined such that you can have an “Active colitis,” a “Chronic active colitis,” or a “Chronic inactive colitis,” which is also sometimes called “Quiescent colitis.”


### Activity = Neutrophils

#### Cryptitis

PMNs in crypt epithelium


#### Crypt Abscess

PMNs in crypt lumen



### Chronicity

Think of those test tubes being melted (like a by a torch)



### Crypt architectural distortion

- Crypt shortening
- Crypt branching
- Crypt dropout
- Loss of crypt parallelism
- Villiform surface


### Basal lymphoplasmacytosis

- Paneth cell metaplasia and hyperplasia
- Pyloric gland metaplasia
- Lamina propria and submucosal fibrosis

*New onset, untreated IBD (hasn't had time for chronicity to develop)*

**Active Colitis**


~ 1 month untreated



*Typical appearance of active disease*

**Chronic Active Colitis**

*Treatment*



*IBD in recent remission (eventually it can normalize)*

**Chronic inactive (Quiescent) Colitis**

## IBD General Info:

Chronic, idiopathic, relapsing and remitting inflammatory disease of the gastrointestinal tract resulting from inappropriate mucosal immune activation. Thought to involve aberrant immune response with altered intestinal microbiome in genetically susceptible individuals.

More common in industrialized nations, where there are fewer parasites/infections to train/distract the immune system ("Hygiene hypothesis").

## Ulcerative Colitis

Chronic active inflammation in the **rectum** proceeding proximally in **continuous, diffuse** pattern

Typical findings:



**Chronic Active Colitis** limited to mucosa and superficial submucosa with ulceration

Can see deeper inflammation with severe "fulminant" colitis

Can have increased inflammation in cecum near appendiceal orifice ("cecal patch")

Can have inflammation in terminal ileum ("backwash ileitis").

\*\*In Kids, can have upper tract findings, relative rectal sparing, and less initial chronicity.\*\*



## Crohn's Disease

**Patchy Transmural** chronic active inflammation in **any** part of the GI tract

Typical findings:

**Transmural** inflammation

**Skip** areas and **patchy** inflammation, both microscopically and grossly

**Granulomas**



**Ulcers:** superficial aphthous to fissuring

Muscle and nerve hypertrophy

**Pyloric gland metaplasia** (esp. in TI)

Fibrosis and strictures

Fistulas



## Indeterminate Colitis

*aka: IBD, type unclassified*

Approximately 10% of patients unclassifiable, due to the pathologic and clinical overlap between UC and CD. Placeholder term--this is **NOT** a specific entity. Often due to insufficient data or fulminant colitis.

## **Differential Diagnosis:** IBD is a diagnosis of exclusion!

### **Active Colitis**

(aka Acute Self-limited Colitis)

**Causes:** E. Coli, Salmonella, Shigella, Campylobacter, Viruses

E. coli O157:H7 → ischemic changes

**Looks similar:** Some medications (e.g., NSAIDs, Checkpoint inhibitors), New onset IBD

### **Neutrophilic Cryptitis** **But, Chronicity ABSENT**

Relatively more neutrophils in superficial lamina propria  
(away from crypts)

Crypt abscesses

Hemorrhage, edema

Possible erosions

### **Focal Active Colitis**

#### **FOCAL Neutrophilic Cryptitis** **Chronicity ABSENT**

**Causes:**

NSAIDS → + Increased apoptoses, ischemic-like changes

Bowel preparation artifact → + Increased apoptoses, edema, mucin depletion

Early infection → Days 0-4 after onset

Ischemic changes → often with lamina propria hyalinization, crypt withering

### **Microscopic Colitis**

#### **Increased Intraepithelial Lymphocytes (IELs)** Neutrophils rare to absent

#### **Lymphocytic Colitis**

IEL ≥20/100 surface epithelial cells

Normal architecture

Chronic inflammation in lamina propria  
(usu. superficial)

#### **Collagenous Colitis**

IEL >10-20/100 surface epithelial cells

Increased Subepithelial Collagen

Entraps capillaries and lymphocytes

Highlighted by Trichrome stain

### **Additional Diagnoses to Consider:**

**Top infectious causes of colitis in patients with IBD:** CMV and C. difficile → always get CMV IHC in a patient with refractory IBD now with severe disease on treatment (esp. steroids)

**If Older** → Especially rule out Medication-effect and Diverticular disease

**Common Drug pattern:** Intraepithelial lymphocytes, relatively preserved crypt architecture, apoptoses, with some neutrophils

**Ischemic colitis** → Hyalinized lamina propria, withered crypts, minimal inflammation

**Radiation colitis** → Ischemic changes, Atypical stromal cells, Telangiectatic blood vessels

**Diverticular disease-associated colitis** → In colonic segment with diverticulosis

**Diversion colitis** → Colon isolated from fecal stream, Follicular lymphoid hyperplasia

**Prolapse** → Fibromuscular hyperplasia, Angulated diamond-shaped crypts

**Vasculitis** → Inflammatory destruction of vessels, Fibrinoid necrosis

**Eosinophilic/Allergic Colitis** → >60 Eos/10 HPF, Few PMNs, Absent chronicity

**STD Proctitis** → Often chlamydia or syphilis due to anal receptive intercourse. Lots of ulceration, plasma cells, and histiocytes. Confined to rectum.

## Medical Management

Usually 2 phases: 1) **Induction** (to induce remission) and 2) **Maintenance** (to maintain remission)  
These may use same or different medications/dosages.

Typical management previously involved “**Step-up** therapy,” where you start with a mild drug (e.g., mesalamine) and only move up to a more powerful drug if they “fail” that drug. However, recent clinical trials have shown better complication-free survival with a “**Top down**” model where you start with a more powerful medication (e.g., monoclonal antibody).

**Mesalamine** (5-ASA) – mechanisms of action unknown. Low activity. Usually used orally or rectally for mild UC.

**Sulfasalazine** – like 5-ASA (mechanism of action unknown). Usually used for mild ileocolic CD.

**Budesonide** – steroid taken orally with little system effect (mainly works on GI tract).

**Prednisone** – oral steroid often used to induce remission in active IBD. Long-term use limited due to side effects. Use in both CD and UC.

**Azathioprine/6-Mercaptopurine** – Thiopurines, inhibit DNA synthesis, thereby reducing WBC production and inflammation. Risk of lymphoma. Used in both CD and UC.

**Tofacitinib** (Xeljanz) – janus kinase (JAK) inhibitor. Currently only used in UC. Oral pill. Powerful.

### Monoclonal antibodies:

**Adalimumab** (Humira) – recognizes TNF $\alpha$ . Used in both CD and UC.

**Infliximab** (Remicade) – recognizes TNF $\alpha$ . Used in both CD and UC.

**Vedolizumab** (Entyvio) – recognizes  $\alpha 4\beta 7$  (gut-specific) integrin, inhibiting diapedesis. Used in both CD and UC, but likely better for UC. Very few side-effects as gut-specific.

**Ustekinumab** (Stelara) – recognizes interleukin (IL) 12 and 23. Used in CD.

**Risankizumab** (Skyrizi) – recognizes interleukin IL-23. Used in CD.

## Cancer Risk and Screening

Inflammation  $\rightarrow$  DNA oxidation/damage  $\rightarrow$  Cancer  
Risk proportional to severity/duration of inflammation.

### **Screening recommendations:**

First 8-10 yrs after diagnosis  $\rightarrow$  No increased screening (not enough time for carcinogenesis)

Years 10-20  $\rightarrow$  Every 1-3 yrs (shorter interval with worse, esp. if PSC)

Years 20 onward  $\rightarrow$  1-2 yrs

### Cancer Risk:

Ulcerative colitis = ~2.4 fold risk (**~ 2x risk**)

Crohn's Disease = ~1.9 fold risk

## Treatment of Dysplasia

With modern techniques, including high-definition and chromoendoscopy, most dysplasia is visible. As such, it can often be completely resected endoscopically.

Once a dysplastic lesion has been resected, in the absence of surrounding dysplasia, ongoing meticulous colonoscopic surveillance is appropriate.

Proctocolectomy is only recommended for dysplasia if endoscopic resection is not possible, or if nonvisible high-grade dysplasia or adenocarcinoma is found.

## Pre-malignant and Malignant lesions in IBD:


Generally, follows stepwise progression of: Non-neoplastic → Low-grade dysplasia → High-grade dysplasia → Adenocarcinoma. However, there are cases where it appears to go from low-grade (or even normal appearing) to adenocarcinoma very quickly or directly.

## Conventional Dysplasia (look like usual colon adenomas):

### Indefinite for Dysplasia

Unable to classify as definitely reactive or dysplastic.  
Often atypia in setting of severe inflammation or ulceration.  
Sometimes surface not present for evaluation.

**Management:** Treat active disease and repeat biopsy in 3-12 months.




### Low-Grade Dysplasia

Looks like a sporadic **Adenoma**.  
Enlarged, hyperchromatic, smooth, "pencil" nuclei.  
Pseudostratified nuclei with maintained basal orientation.  
Higher N:C ratios; **Little to no surface maturation.**  
Often abrupt transition (corresponding with clone)  
Prominent apoptoses.

**Molecular:** IBD-associated dysplasia show more copy number aberrations and aneuploidy than sporadic adenomas. TP53 mutations are very frequently present early. Possibly reflecting a faster progression toward cancer.

**Management:** Complete endoscopic resection if visible.  
Otherwise proctocolectomy ± IPAA to exclude cancer.




*Hint: Try using a lymphocyte as what is "normochromatic"*

### High-Grade Dysplasia

Enlarged, hyperchromatic, pleomorphic nuclei.  
Often plumper than LGD.  
Irregular nuclear contours. Prominent nucleoli.

#### **Loss of nuclear polarity.**

Complex architecture: Cribriforming, crypt branching/budding.



**Immunohistochemistry in IBD dysplasia:** P53 staining often highlights both grades  
Dysplasia → Strong P53 staining (or null). *Some* authors require abnormal p53 *at the surface*, while others just want it to be significantly increased compared to the background colon.

Negative/Indefinite for dysplasia if weak/wild-type staining

SATB2 is frequently lost in IBD dysplasia also, but this is used less often as a marker.

H&E is still the gold standard though, so only do it on cases that are equivocal!

# **Nonconventional lesions:** (doesn't look like usual colon adenomas)


## **Serrated Epithelial Change**

### ***Controversial diagnosis, with differing criteria***

**Original Hopkins Criteria:** Serrations at top and bottom of crypts. Distorted crypt architecture where some crypts do not reach the muscularis mucosae (unlike SSL). Normal nuclei. Goblet cell-rich epithelium.

**UCSF Criteria:** Hyperplastic polyp (HP)-like mucosal change without morphologic evidence of dysplasia detected on random biopsy

**Controversial** risk of CRC. Many studies show increased risk of dysplasia/carcinoma. Essentially, treat as "indefinite."



*Note: The colon in IBD patients can show surface serrations/hyperplasia, particularly in the distal colon, making this diagnosis especially hard and controversial.*

## **Non-Conventional Dysplasia**

Often present with conventional dysplasia. More common on left side as polypoid mass.

**Hypermucinous**—Villous architecture with prominent cytoplasmic mucin.


**Traditional Serrated Adenoma (TSA)-like**

**Sessile serrated lesion (SSL)-like**

**Paneth cell differentiation**

**Goblet cell deficient**—absence of goblet cells

**"Terminal epithelial differentiation,"** TED, or "Crypt cell dysplasia," CCD —flat lesions, round to oval hyperchromatic nuclei. Can be just in crypts.



**Helpful tip:** Must see **cytologic atypia (hyperchromatic, crowded nuclei)**.

In uncertain cases, get p53 IHC. If not altered → better to hedge as "indeterminate for dysplasia" and state change type (e.g., "Hypermucinous epithelium, Indefinite for dysplasia")

## **Adenocarcinoma**

Invasive through basement membrane:

- Infiltrating glands/cells
- Broad, expansive confluent growth of glands

Compared to Sporadic, IBD-associated CRC is:

- More often multifocal (field defect)
- More often higher grade
- More often advances stage
- More often signet-ring or mucinous

Unique variant:

**Low-grade tubuloglandular adenocarcinoma**—very bland small to medium-sized round glands that invade with little desmoplastic stroma. Often CK7 (+). Loss of SATB2. Frequent IDH1 mutations.

